

# Neuronal plasticity and antidepressant actions

Eero Castrén<sup>1</sup> and René Hen<sup>2</sup>

<sup>1</sup> Neuroscience Center, University of Helsinki, P.O. Box 56, 00014 Helsinki, Finland

<sup>2</sup> Departments of Neuroscience and Psychiatry, Columbia University, New York, NY 10032, USA

**Antidepressant treatments enhance plasticity and increase neurogenesis in the adult brain, but it has been unclear how these effects influence mood. We propose that, like environmental enrichment and exercise, antidepressant treatments enhance adaptability by increasing structural variability within the nervous system at many levels, from proliferating precursors to immature synaptic contacts. Conversely, sensory deprivation and chronic stress reduce this structural variability. Activity-dependent competition within the mood-related circuits, guided by rehabilitation, then selects for the survival and stabilization of those structures that best represent the internal or external milieu. Increased variability together with competition-mediated selection facilitates normal function, such as pattern separation within the dentate gyrus and other mood-related circuits, thereby enhancing adaptability toward novel experiences.**

## Neuronal plasticity: growth and change

The extrinsic and intrinsic milieu becomes represented in the structure and function of neuronal networks during development through neuronal plasticity. In adulthood, this representation is continuously optimized and modulated through plasticity and learning [1]. Plasticity occurs at several levels, from neurogenesis to the adjustment of synaptic weights, and modulates both the structure and function of neuronal networks (Figure 1). Changes in function of the nervous system are essentially always based on a structural change at some level (neuronal, synaptic, protein, or genomic structure) and it is therefore difficult to make clear distinctions between functional and structural plasticity.

Plasticity can be conceptualized as two distinct processes. First, structural variability is generated through the overproduction of immature neuronal structures, which occurs at many levels from newborn neurons to the outgrowth of filopodia (Figure 1). Second, selective stabilization among the overproduced structures retains those that best represent the internal or external milieu [2]. Structural variability may be stochastic, although it may also be genetically tuned, but selection is an active process driven by neuronal activity that reflects both extrinsic and intrinsic stimuli. Activity-dependent selection guides

the stabilization of functionally relevant neurons and connections by utilizing genes involved in survival and synaptogenesis, or it eliminates weakly or incoherently active structures through the use of genes that underlie apoptosis and pruning [1,3]. The elimination of weakly active neurons and connections is critical for optimizing the signal-to-noise ratio in neuronal networks. Neuronal plasticity can be compared to auditions for a Broadway show: if many candidates audition, the production team can select an optimal performer for each role (and send those not suitable back home), but if only a few people show up, almost all have to be utilized regardless of their talent.


During the past few years, neuronal plasticity, and in particular adult neurogenesis, has been implicated in the beneficial effects of antidepressant drugs and electroconvulsive shock (ECS) treatment [4–7]. However, it has remained unclear how plasticity and neurogenesis impact mood and anxiety-related behaviors.

Here, we provide a framework for how chronic antidepressant drug treatment and ECS might utilize neurogenesis and other forms of neuronal plasticity to influence mood. We argue that antidepressants, environmental enrichment, and exercise increase structural variability at various levels in the nervous system, thereby offering more substrates for the selection process (Figure 1) [7]. Conversely, sensory deprivation and chronic stress would reduce variability and impair adaptability. Whereas acute stress may promote variability and adaptation, chronic stress is considered maladaptive and is associated with a loss of neurons and synapses [8,9]. Furthermore, stress may increase activity in certain brain regions, such as the amygdala and the mesolimbic dopaminergic system, leading to hypertrophy of these structures.

We argue that chronic antidepressant treatments and ECS act – at least in part – by utilizing a similar mechanism to increase adaptability and facilitate structural and functional reorganization in neuronal networks that have evolved to boost the effects of environmental enrichment, although the cellular and molecular mechanisms may be different. We further suggest that the effects of enrichment and antidepressant treatments that increase plasticity take place at many levels within neuronal networks (Figure 1). Because experience-guided selection constantly eliminates inactive structures, an increase or decrease in variability does not need to influence the mean number of these structures in the brain (like counting the number of actors of the final cast does not tell you how many have auditioned), which makes detecting such a change

Corresponding authors: Castrén, E. (eero.castrén@helsinki.fi); Hen, R. (rh95@columbia.edu).

Keywords: antidepressants; enrichment; stress; neurogenesis; synaptogenesis; plasticity.



**Figure 1.** Proposed model for the levels of neuronal plasticity in antidepressant activity. Neuronal plasticity acts at different structural levels and bidirectionally to influence variability and selection within neuronal networks. Increased formation of structures at any of these levels by, for example, environmental enrichment (EE) or chronic antidepressant treatment generates variability and promotes competition between similar structures for stabilization, thereby enhancing adaptability, even if the total number of structures is not increased due to the simultaneous increase in structural elimination. **(a)** Neurogenesis and selective apoptosis. Increased precursor proliferation in the dentate gyrus, induced by antidepressants and EE, leads to the increased survival of newborn neurons that successfully mediate activity within the hippocampal circuitry. Neurons that fail to functionally integrate into the hippocampal circuitry are eliminated through apoptosis. **(b)** Arborization and pruning of axonal and dendritic branches. The increased dynamics of nascent branches promotes the stabilization of branches containing synapses that successfully represent environmental conditions, whereas arbors without active synapses remain short lived and are pruned. **(c)** Synaptogenesis and synaptic elimination. Immature 'trial synapses' between two neurons are initiated by filopodial extension from pre- or postsynaptic sites. Synapses that are successfully activated during the trial period are preferentially selected for stabilization, whereas contacts that fail to mediate activity collapse and are eliminated. **(d)** Plastic regulation of synaptic strength. Information transfer through active synapses is potentiated through the process of long-term potentiation (LTP), whereas inactive or inappropriately active synapses are suppressed through long-term depression (LTD). **(e)** Environmental activity regulates the transcription and translation of effector genes involved in neuronal plasticity through transcriptional control and epigenetic mechanisms, such as remodeling of chromatin structure from a closed to an open state.

in variability difficult. Therefore, it is currently unclear whether the effects of antidepressants are confined to particular brain areas or whether they occur more ubiquitously throughout the brain. Nevertheless, tougher competition between variable structures increases adaptability toward changes in the external or internal milieu, analogous to the way increased genetic variability contributes to the survival of a species facing a changing environment in the context of evolution.

### Neurogenesis in the dentate gyrus

Adult neurogenesis is an exceptional form of plasticity in which entire neurons are generated and selected for survival in only a very few regions of the adult mammalian brain [10]. The two main areas where neurogenesis occurs are the subventricular zone that lines the ventricles and gives rise to neuronal precursors that migrate toward the olfactory bulb and the subgranular zone that lines the dentate gyrus (DG) of the hippocampus and gives rise to DG granule cells (Figure 2) [10].

Neurogenesis is a plastic process that is regulated by environmental factors. In the hippocampus, various stages of neurogenesis are stimulated by enriched environments, exercise, learning, and antidepressant drugs, and they are inhibited by chronic stress and aging [5,6,11]. Enrichment, antidepressants, and ECS stimulate the proliferation of neural stem cells, their differentiation into neurons, and the survival of the resulting young neurons (Figure 2) [4,12]. Conversely, both acute and chronic stress decrease the proliferation and survival of newborn neurons in several species, including nonhuman primates [13,14], although increased neurogenesis has not been confirmed in all stress studies [13,14]. Newborn neurons have also been implicated in the adaptation to stress by buffering against it [15]. Furthermore, sensory deprivation decreases proliferation, the choice of neuronal fate, and the survival of the young neurons [16]. These observations raise the possibility that changes in neurogenesis allow for better adaptation to a changing environment either instructively by encouraging adaptive behavior [17] or permissively by increasing variability that is then utilized in experience-dependent selection. However, it is also possible that changes in neurogenesis are merely a consequence of these environmental changes. By reviewing the proposed functions of adult-born neurons, we can attempt to resolve these questions.


### Neurogenesis, pattern separation, and generalization

Recent functional studies of adult-born dentate granule neurons have focused on their potential role in pattern separation because increasing evidence from electrophysiological studies indicates that the DG is involved in pattern separation [18]. It has been proposed that pattern separation enables the processing of similar experiences as distinct memories and is critical for memory formation, although direct evidence for this mechanism is still missing. For example, pattern separation may enable us to remember two distinct beach vacations or where we last parked our car, even though the parking garage and general context may be the same each morning.

In numerous rodent models, both loss- and gain-of-function studies show that increases in neurogenesis improve and decreases in neurogenesis impair pattern separation [19,20]. Furthermore, some evidence exists indicating that young neurons are involved in discriminating between complex odor mixtures in the olfactory bulb, which may involve a process similar to pattern separation [21].

Pattern separation appears to be impaired both during normal aging and in individuals with mild cognitive impairment [22]. In addition, functional imaging studies have identified abnormal activity in the dentate gyrus and CA3 of individuals with age-related memory impairments when they perform a pattern separation task [23].

In the psychiatric literature, the term pattern separation is rarely used because it refers to a cognitive process that is usually not tested in psychiatric patients. However, generalization is a phenotype that is often associated with anxiety and mood disorders [24]. Generalization can be defined as our tendency to lump together similar experiences, particularly when they have a strong emotional



**Figure 2.** Effects of mood and environment on pattern separation and generalization. Information processing within the trisynaptic loop of the hippocampus [dentate gyrus (DG) → CA3 → CA1] is required for discrimination between similar contexts. This process is termed pattern separation and is modulated by adult neurogenesis within the DG (red boxes). Through the differing connectivities of the dorsal and ventral hippocampus (association cortices versus amygdala, bed nucleus of stria terminalis (BNST), prefrontal cortex, and hypothalamus), both neutral and emotionally charged contextual information is processed and discriminated. When adult neurogenesis is reduced or blocked by stress, aging, or experimental manipulations (such as X-ray treatment), discrimination is impaired, leading to generalization [17,19]. By contrast, increased neurogenesis, promoted by environmental enrichment, antidepressant treatment, or genetic manipulations (e.g., Bax inhibition), improves pattern separation and discrimination [19,36]. Abbreviations: Amyg, amygdala; PFC, prefrontal cortex; Hypoth, hypothalamus; Antidep, antidepressant treatment.

content. For example, the sight of a plane flying over a skyscraper may remind us of 9/11. Generalization may therefore be considered the opposite of pattern separation. Generalization is a double-edged sword. In small doses, it is clearly protective to avoid similar dangerous situations – for example, if you were hit by a car when you crossed a dangerous intersection, a little generalization will enable you to be careful each time you cross this intersection; however, too much generalization may result in the fear of crossing any street, which would clearly be maladaptive. Like pattern separation, generalization may be critical for memory formation because it may allow us to link similar memories rather than store them in unrelated categories. In the cognitive domain, the process that allows generalization is pattern completion, which has been proposed to take place in CA3 [25]. Thus, when new memories are encoded two processes are simultaneously at work: pattern separation to disambiguate similar situations and pattern completion (or generalization) to link similar or related events, particularly when they have a strong valence. There is indeed evidence that, during memory encoding, a balance between separation and completion allows similar experiences to be stored either together or separately. For example, from an evolutionary point of view, it is clearly important to store aversive memories together because any experience that is similar to a traumatic memory should be avoided.

Therefore, the DG appears to function in different regimes in different environmental situations. In a safe and rich environment, high pattern separation is advantageous because it increases the ability to discriminate between similar experiences, which may be optimal for a situation of high exploratory activity aimed at locating food

and mates. By contrast, in a dangerous environment low pattern separation or generalization may be preferable because it will increase fear and promote avoidance rather than exploration [17,19].

The idea that variable levels of pattern separation have adaptive value raises the possibility that variable levels of neurogenesis may promote adaptation to a changing environment. The mechanisms through which increased neurogenesis contributes to pattern separation remain unclear (Box 1), but the fact that a major period of cell death occurs around the time that the neurons have established both pre- and postsynaptic contacts suggests that the ability to contribute to the activity of the DG is critical. This further suggests that adult neurogenesis may select the surviving neurons using principles similar to those that have been thoroughly investigated in the context of peripheral nervous system development. During early development, sensory and sympathetic neurons are produced in excess and when their axons reach their target tissues, newborn neurons compete for access to a neurotrophic factor produced by the target cells. Neurons that establish an optimal connection with the target cells receive a sufficient amount of the trophic factor and survive, whereas those that fail to optimally innervate the target are eliminated by programmed cell death [26,27].

It is conceivable that a similar principle of redundancy and competition governs the selection of surviving neurons in the mammalian DG, where neurogenesis continues into adulthood and where neuronal production, selection, and elimination are continuously taking place. Although the hypothetical target-derived neurotrophic factor for newborn DG neurons has not been identified, this factor should be regulated and released in an activity-dependent manner

**Box 1. Outstanding questions**

The hypothesis that increase in structural variability induced by enrichment and antidepressant treatments promotes adaptability after activity-dependent selection is largely based on information derived from experiments performed in the peripheral nervous system and in primary sensory systems. Several important questions remain to be addressed before these principles can be extended to the higher cortical regions, such as those involved in the regulation of mood. These questions include:

- Do antidepressants and enrichment promote synapse turnover in higher brain regions, such as the PFC or the hippocampal CA1 area?
- If enrichment and antidepressant treatments enhance variability, does this promote adaptability in mood-related behavior?
- Are the effects of antidepressants confined to particular brain areas or do they occur more ubiquitously throughout the brain?
- What mechanisms select newborn neurons for survival?
- Enrichment and antidepressants reactivate critical-period-like plasticity in the adult visual cortex in rodents, but do they produce similar reactivation in the human brain?

[28]. Indeed, brain-derived neurotrophic factor (BDNF), one of the prime candidates for this function, is regulated by neuronal activity [29]. We propose that, when environmental conditions are changing, a newborn DG neuron with a slightly different activity pattern, reflecting different patterns of sensory input, has a higher probability to be selected for survival than newborn cells displaying activities very similar to the neurons that already exist in the mature DG. This idea is consistent with the increased survival of newborn neurons in an enriched environment. At least in the case of antidepressant treatment, there is evidence that increased survival of newborn neurons is balanced by an increase in the total number of apoptotic neurons within the DG, suggesting that although the newborn neurons are surviving, older neurons are eliminated [30]. Analogously, a Broadway producer planning for a new show looks forward to a large number of auditionees with diverse talents to replace at least some members of the current cast, even if the current cast was excellent in the previous show. Therefore, when the input to the DG is variable, enhanced neurogenesis may be beneficial because it favors pattern separation.

**Neurogenesis and antidepressant action**

Plasticity that is based on neurogenesis operates on a different time scale than traditional forms of plasticity, such as spine or dendritic rearrangements that can occur much faster than the generation of new neurons. If variable levels of neurogenesis have an adaptive value, it is likely to be in response to long-lasting environmental changes such as those resulting from changing seasons. For example, neurogenesis varies in the hippocampus of birds that store food in hidden caches for retrieval in the winter. In these birds, neurogenesis is highest in the fall and winter when they hide and retrieve their food [31]. Similarly, treatments that increase neurogenesis are unlikely to produce a rapid behavioral response. This may be one reason why antidepressants have a delayed onset of therapeutic effect.

Evidence from animal models of anxiety and depression indicate that neurogenesis is necessary for some but not all effects of antidepressants [32,33]. Given the role of neurogenesis in pattern separation, we hypothesize that an

improvement in pattern separation, particularly for situations and contexts that are emotionally charged, will impact mood and anxiety-related behaviors. Such an effect may be achieved by connections between the ventral hippocampus and the limbic system [34]. Unlike the dorsal part of the hippocampus, which sends projections primarily to association cortices, the ventral hippocampus also sends projections to the amygdala, the bed nucleus of the stria terminalis, the hypothalamus, and the prefrontal cortex (PFC). In the PFC, projections from the ventral hippocampus have recently been shown to activate neurons that fire in response to anxiety-related modalities [35]. It is therefore possible that a particular context acquires a valence by virtue of the connections between the ventral hippocampus and the limbic system.

The process of neurogenesis may be harnessed to improve cognition and mood. A recent study demonstrated that inhibition of apoptotic cell death among the progeny of hippocampal neural stem cells significantly increases neurogenesis and improves pattern separation [36]. Interestingly, this genetic manipulation had no impact on anxiety-related behaviors unless it was combined with exercise, which is consistent with the idea that enhanced survival benefits from the increased variability provided by exercise-induced precursor proliferation. Future studies in animal models are needed to investigate whether the combination of strategies aimed at stimulating neurogenesis (by inhibiting cell death or other means), together with behavioral enrichment or exercise, will result in antidepressant or anxiolytic-like effects.

**Plasticity outside the DG**

Increasing evidence suggests that enhanced neuronal plasticity induced by enrichment and chronic antidepressant treatment may not be restricted to neurogenesis (Figure 1). Furthermore, the mechanisms that mediate the effects of these treatments at smaller structural scales may be conceptually similar to those reviewed above for neurogenesis. However, technical limitations have hampered the recognition of these effects; it is difficult to detect changes in the turnover of dendritic branches and spines when there are simultaneous changes in the rates at which these structures are produced and retracted without any change in the net number of structures. Therefore, it is currently unclear whether the effects of antidepressants are confined to mood-related circuits or are more widespread (Box 1). Nevertheless, at least in primary sensory areas, a change in turnover has significant functional consequences [37,38]. Developments in intravital microscopy of behaving animals is now circumventing these technical difficulties and an increasing number of reports have focused on the dynamic effects of environmental manipulation on the structure of dendrites and axons [39,40].

**Axonal and dendritic branches**

It has been proposed that the construction of neural circuits proceeds through concurrent and nearly balanced growth and retraction of axonal and dendritic branches [3,41–43]. Nascent dendritic branches have been proposed to produce ‘trial synapses’, and only those trial synapses that receive appropriate synaptic input are preserved and



the corresponding dendritic branch stabilized [3]. Therefore, activity-dependent synapse stabilization appears to direct axonal and dendritic arbor selection and elimination (Figure 1).

There is evidence that antidepressant treatment and ECS increase the variability and turnover of the branches of dendrites and axons (Figure 1). Chronic fluoxetine administration simultaneously increases the elongation and retraction of branch tips in the mouse visual cortex [40] and ECS promotes axonal sprouting in the hippocampus [44], consistent with the idea that antidepressants increase variability in branch dynamics. Conversely, chronic mild stress has been shown to reduce the volume as well as the length and branching of apical dendrites within the DG, the CA3 area, and the PFC in rats [45]. All of these effects were reversed by antidepressant treatment [45].

### Synaptic connections

During early postnatal development, the density of synapses in the human cortex exceeds that found in the adult brain by about twofold [46,47]. Although brain growth contributes to reduced spine density, it is thought that a net loss of synapses brings the synaptic density to the adult level at adolescence. At least in the human PFC, synaptic pruning continues well into the third decade [47]. It has been proposed that synaptic activity selects from the over-produced synapses those that are stabilized [2]. This activity-dependent process ensures that only synapses that optimally represent external or internal input are retained and that those mediating random noise are eliminated [3,42].

Even in adulthood, synaptogenesis continues at a lower level, but if synaptic elimination occurs at a matching rate, the net number of synapses remains stable. Evidence from sensory cortices suggests that a simultaneous increase in spine formation and retraction increases adaptability by making a higher number of trial contacts available for selection (Figure 1) [40,48,49], as when a larger number of auditionees helps in selecting the optimal cast for a show. In higher cortical areas, evidence for a correlation between increased synapse turnover and improved function is lacking, due at least in part to technical difficulties (Box 1). It is possible that currently available methods underestimate the dynamic changes in the turnover of synaptic contacts that may take place after environmental changes or antidepressant drug administration (Figure 1c).

ECS increases the number of synapses in the hippocampus, but antidepressant treatment has only a minor effect on the net number of dendritic spines in the hippocampal CA1 area [50,51]. However, when the number of spines and synapses is abnormally downregulated by stress [52] or ovariectomy [53], the increasing effect of antidepressants on spine number becomes unmasked and fluoxetine treatment increases spine number back to the baseline level. This suggests that, in the normal hippocampus, fluoxetine might simultaneously increase spine formation and elimination, thereby having only a minor effect on the net number of spines, but fluoxetine may produce this effect also indirectly through other mechanisms. A recent study that observed dendritic contacts repeatedly using two-photon microscopy in the mouse visual cortex reported

that chronic fluoxetine treatment simultaneously increased the elongation and retraction of dendritic branch tips [40].

Chronic stress and long-term glucocorticoid treatment lead to the loss of dendritic spines and synaptic contacts in the hippocampus and the PFC [8,54–56]. Chronic mild stress also increases the number of immature spines at the expense of mushroom-like mature spines in the apical dendrites of pyramidal neurons in the hippocampus and PFC and these changes are largely reversible by antidepressant drug treatment [45]. A recent study reported that short-term glucocorticoid treatment increased spine dynamics in the mouse somatosensory cortex by simultaneously increasing spine formation and retraction, and that inhibition of endogenous glucocorticoids reduces spine dynamics [9]. Consistent with earlier findings, long-term glucocorticoid treatment increased net spine elimination.

### Plasticity of synaptic strength

In addition to synapse number, synaptic strength is also dynamically regulated by environmental experiences, including enriched environment, exercise, and antidepressant drugs. Chronic fluoxetine administration increases long-term potentiation (LTP) in the DG elicited in the absence of GABA<sub>A</sub> receptor (GABA<sub>A</sub>R) inhibitors and this effect depends on the newborn neurons [12]. In the presence of GABA<sub>A</sub>R inhibitors, DG LTP is reduced, perhaps due to occlusion, and long-term depression (LTD) is enhanced [12,57,58]. Enrichment and fluoxetine enable LTP in the adult rat visual cortex [48,59] and a similar effect of fluoxetine treatment on LTP was observed in the murine amygdala [60]. These findings may be related to the ‘dematuration’ process observed after chronic fluoxetine administration in the dentate granule neurons [58], indicating that antidepressant treatment reactivated a juvenile-like plasticity in the brain [48,60]. Enriched environment and perhaps also fluoxetine treatment during early life accelerate cortical maturation [61–63]. Conversely, chronic mild stress facilitates LTD in the CA1 area and chronic antidepressant treatment blocked this LTD facilitation and enhanced LTP [64]. Thus, chronic antidepressant treatment and enriched environment may increase synaptic plasticity in several brain areas (Figure 1), which may be consistent with the increased dendritic spine dynamics and turnover induced by antidepressant treatment [40].

### Genomic plasticity


The regulated expression and translation of specific genes by experience-dependent neuronal activity is a critical mechanism of neuronal plasticity [65–67]. Neuronal activity can influence gene expression by activating transcription factors or by inducing epigenetic changes in chromatin structure or DNA methylation [68–73] (Figure 1). Mutations in several genes regulated by activity are associated with neurodevelopmental disorders, which underlines the critical importance of this process for proper network connectivity [65,66].

BDNF is a critical mediator of neuronal activity and synaptic structure [29,74] and its production and release are regulated by neuronal activity [29]. BDNF signaling

through TrkB receptors promotes the survival of newborn neurons in the DG [30,75,76], enhances the outgrowth of axons and dendrites [43,77], stabilizes synapses, and promotes synaptic transmission [78,79]. The effects of enrichment and antidepressant drugs are at least partially mediated by BDNF signaling [30,48,76,80,81], although in brain areas that are activated by aversive stimuli, BDNF has pro-depressive effects [82]. Thus, the activity-dependent regulation of BDNF is a critical molecular mediator through which experience-dependent plasticity is translated into structural and functional changes in neuronal networks. If this were a Broadway show, BDNF

would be a producer that selects actors and actresses for the cast from among the auditioning candidates.

Recent experimental and theoretical work suggests that environmental conditions may have relatively small or variable effects on the expression of individual genes, but they reliably increase the large-scale variability in gene expression, or the 'genomic tone' [83,84]. High local variability in the DNA methylation rate without any change in the mean methylation level was recently discovered within variably methylated regions of genomes from several species [85] and may provide a mechanism for changes in the genomic tone [84]. This altered variation



**Figure 3.** Plasticity of neuronal networks in mammalian primary visual cortex. **(a)** Soon after eye-opening, inputs that mediate visual information from the left (blue) and right eye (red) diffusely innervate the entire layer IV of the primary visual cortex. **(b)** During the postnatal critical period (CP), activity-dependent competition leads to the segregation of inputs from either eye into eye-specific regions, the ocular dominance (OD) columns. As a consequence, at the end of critical period, cortical neurons within a single column in layer IV receive innervation predominantly from a single eye (middle panel). This segregation requires vision-induced neuronal activity such that axonal branches from one eye withdraw from the regions initially dominated by the other eye and elaborate connections in their own territory (lower panel). It should be noted that the OD columns cannot be detected in the primary visual cortex as visible structures, even though they are differentially colored in the figure for the sake of clarity. **(c)** Development of a normal network requires balanced use of both eyes. If one eye is deprived of vision during the critical period, the inputs mediating information from it lose in activity-dependent competition and withdraw [38]. The inputs mediating visual information from the open eye are active and overcome most of the visual cortex. If the vision of the deprived eye is not corrected and encouraged during the critical period, the network guided by only one open eye remains permanent even if the deprived eye is opened in adulthood. **(d)** Fluoxetine treatment (Flu) [48] or exposure to an enriched environment in adulthood [59] reopens the critical-period-like plasticity. This enables a reorganization of the network, if the deprived eye is opened in adulthood and encouraged by temporary patching of the previously open eye. Under these conditions, the now more active deprived eye can regain territories within layer IV, which gradually leads to a normal network guided by two open eyes. Note that under all of these conditions, the net number of synapses remains constant (lower panel); only the source of visual information varies.

in methylation and gene expression without any change in the mean resembles the increase in structural variation and turnover in neurons and synapses discussed above, but it is unclear whether and to what extent the genome-level variation is causally related to variation in network dynamics. However, variability in gene expression levels does correlate with behavior [83]. Intriguingly, genes adjacent to the variably methylated regions have often been found to be functionally related to brain development and plasticity, even when the tissue analyzed was liver [85]. This suggests that variably methylated regions are associated with many genes involved in neuronal plasticity, which may induce variability in their expression levels throughout the organism.

### The visual cortex as an example of experience-dependent plasticity

Critical-period plasticity in the mammalian visual cortex is a well-characterized model for cortical development and plasticity [37,38,86,87] (Figure 3). It is widely thought that similar processes govern the development and tuning of neuronal connectivity in other cortical areas [1,87]. Recent studies have revealed that critical-period-like plasticity can be reactivated in the adult visual cortex by numerous treatments, including enrichment and chronic fluoxetine treatment (Figure 3d) [40,48,59,62,88,89].

Reactivation of developmental plasticity in the adult brain is apparently not restricted to the visual cortex. Chronic fluoxetine treatment induces dematuration of neurons in the mouse DG that extends to the already matured granule neurons [58]. A recent study used the fear-conditioning paradigm to show that chronic fluoxetine treatment increases neuronal plasticity in the amygdala and leads to long-term removal of the conditioned fear response when fluoxetine treatment is combined with extinction training; neither fluoxetine treatment nor extinction training alone produced long-term fear removal [60]. These findings demonstrate that enriched environment or fluoxetine treatment in adult animals reactivates a critical-period-like plasticity that facilitates the reorganization and functional recovery of a network miswired during development.

### Concluding remarks

Taken together, the evidence above suggests that plasticity-inducing treatments such as exercise, enrichment, and chronic antidepressant treatment increase variability at several structural levels of the nervous system. At all levels, new neurons or new synapses are produced in excess and compete for survival or stabilization. Neurons and synapses that contribute to the activity within neuronal networks are selected for survival and this experience-dependent competition guides the network structure to better represent the external and internal milieu. These data suggest that the combination of enrichment or fluoxetine treatment with rehabilitation could be useful in numerous conditions where the activation of adult plasticity would be desired. Indeed, recent studies have shown the usefulness of antidepressant treatment for recovery from stroke in humans [90] as well as in animal models of Alzheimer's disease [91,92] and traumatic brain injury

[93]. For the treatment of depression, the data reviewed above suggest that antidepressant treatment is effective only when combined with rehabilitation, such as psychotherapy. Clinical trials testing the necessity of rehabilitation in the antidepressant effect should be designed and the combination of antidepressants and psychotherapy, which is recommended by treatment guidelines but too often not followed in clinical practice, should be promoted.

### Acknowledgments

The authors thank Sarah Mack for drawing Figure 1. This manuscript was written during the sabbatical of E.C. at the laboratory of R.H., supported by the Senior Investigator Grant of the Academy of Finland and the Schaefer Scholarship of Columbia University. Original work was supported by the Sigrid Jusélius Foundation, the Academy of Finland Center of Excellence program (for E.C.), the National Institute for Mental Health, the New York Stem Cell Initiative, and Hope for Depression Research Foundation (for R.H.).

### Disclaimer statement

E.C. is cofounder of and advisor to Hermo Pharma, which is running a clinical trial on the use of fluoxetine for amblyopia in adult humans. R.H. is a member of the scientific advisory board of Lundbeck and Roche.

### References

- Katz, L.C. and Shatz, C.J. (1996) Synaptic activity and the construction of cortical circuits. *Science* 274, 1133–1138
- Changeux, J.P. and Danchin, A. (1976) Selective stabilisation of developing synapses as a mechanism for the specification of neuronal networks. *Nature* 264, 705–712
- Hua, J.Y. and Smith, S.J. (2004) Neural activity and the dynamics of central nervous system development. *Nat. Neurosci.* 7, 327–332
- Sahay, A. and Hen, R. (2007) Adult hippocampal neurogenesis in depression. *Nat. Neurosci.* 10, 1110–1115
- Warner-Schmidt, J.L. and Duman, R.S. (2006) Hippocampal neurogenesis: opposing effects of stress and antidepressant treatment. *Hippocampus* 16, 239–249
- David, D.J. et al. (2010) Implications of the functional integration of adult-born hippocampal neurons in anxiety-depression disorders. *Neuroscientist* 16, 578–591
- Castrén, E. (2005) Is mood chemistry? *Nat. Rev. Neurosci.* 6, 241–246
- McEwen, B.S. (1999) Stress and hippocampal plasticity. *Annu. Rev. Neurosci.* 22, 105–122
- Liston, C. and Gan, W.B. (2011) Glucocorticoids are critical regulators of dendritic spine development and plasticity in vivo. *Proc. Natl. Acad. Sci. U.S.A.* 108, 16074–16079
- Ming, G.L. and Song, H. (2011) Adult neurogenesis in the mammalian brain: significant answers and significant questions. *Neuron* 70, 687–702
- van Praag, H. et al. (2000) Neural consequences of environmental enrichment. *Nat. Rev. Neurosci.* 1, 191–198
- Wang, J.W. et al. (2008) Chronic fluoxetine stimulates maturation and synaptic plasticity of adult-born hippocampal granule cells. *J. Neurosci.* 28, 1374–1384
- Schoenfeld, T.J. and Gould, E. (2012) Stress, stress hormones, and adult neurogenesis. *Exp. Neurol.* 233, 12–21
- Vollmayr, B. et al. (2007) Neurogenesis and depression: what animal models tell us about the link. *Eur. Arch. Psychiatry Clin. Neurosci.* 257, 300–303
- Dranovsky, A. and Leonardo, E.D. (2012) Is there a role for young hippocampal neurons in adaptation to stress? *Behav. Brain Res.* 227, 371–375
- Dranovsky, A. et al. (2011) Experience dictates stem cell fate in the adult hippocampus. *Neuron* 70, 908–923
- Glasper, E.R. et al. (2012) Adult neurogenesis: optimizing hippocampal function to suit the environment. *Behav. Brain Res.* 227, 380–383
- Leutgeb, J.K. et al. (2007) Pattern separation in the dentate gyrus and CA3 of the hippocampus. *Science* 315, 961–966
- Sahay, A. et al. (2011) Pattern separation: a common function for new neurons in hippocampus and olfactory bulb. *Neuron* 70, 582–588



- 20 Nakashiba, T. *et al.* (2012) Young dentate granule cells mediate pattern separation, whereas old granule cells facilitate pattern completion. *Cell* 149, 188–201
- 21 Wilson, D.A. (2009) Pattern separation and completion in olfaction. *Ann. N. Y. Acad. Sci.* 1170, 306–312
- 22 Yassa, M.A. *et al.* (2011) Age-related memory deficits linked to circuit-specific disruptions in the hippocampus. *Proc. Natl. Acad. Sci. U.S.A.* 108, 8873–8878
- 23 Yassa, M.A. *et al.* (2010) High-resolution structural and functional MRI of hippocampal CA3 and dentate gyrus in patients with amnesic mild cognitive impairment. *Neuroimage* 51, 1242–1252
- 24 Lissek, S. *et al.* (2010) Overgeneralization of conditioned fear as a pathogenic marker of panic disorder. *Am. J. Psychiatry* 167, 47–55
- 25 Leutgeb, S. and Leutgeb, J.K. (2007) Pattern separation, pattern completion, and new neuronal codes within a continuous CA3 map. *Learn. Mem.* 14, 745–757
- 26 Levi-Montalcini, R. (1987) The nerve growth factor 35 years later. *Science* 237, 1154–1162
- 27 Deppmann, C.D. *et al.* (2008) A model for neuronal competition during development. *Science* 320, 369–373
- 28 Castrén, E. (2004) Neurotrophic effects of antidepressant drugs. *Curr. Opin. Pharmacol.* 4, 58–64
- 29 Thoenen, H. (1995) Neurotrophins and neuronal plasticity. *Science* 270, 593–598
- 30 Sairanen, M. *et al.* (2005) Brain-derived neurotrophic factor and antidepressant drugs have different but coordinated effects on neuronal turnover, proliferation, and survival in the adult dentate gyrus. *J. Neurosci.* 25, 1089–1094
- 31 Barnea, A. and Nottebohm, F. (1994) Seasonal recruitment of hippocampal neurons in adult free-ranging black-capped chickadees. *Proc. Natl. Acad. Sci. U.S.A.* 91, 11217–11221
- 32 Santarelli, L. *et al.* (2003) Requirement of hippocampal neurogenesis for the behavioral effects of antidepressants. *Science* 301, 805–809
- 33 David, D.J. *et al.* (2009) Neurogenesis-dependent and -independent effects of fluoxetine in an animal model of anxiety/depression. *Neuron* 62, 479–493
- 34 Fanselow, M.S. and Dong, H.W. (2010) Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron* 65, 7–19
- 35 Adhikari, A. *et al.* (2010) Synchronized activity between the ventral hippocampus and the medial prefrontal cortex during anxiety. *Neuron* 65, 257–269
- 36 Sahay, A. *et al.* (2011) Increasing adult hippocampal neurogenesis is sufficient to improve pattern separation. *Nature* 472, 466–470
- 37 Hensch, T.K. (2005) Critical period plasticity in local cortical circuits. *Nat. Rev. Neurosci.* 6, 877–888
- 38 Hubel, D.H. *et al.* (1977) Plasticity of ocular dominance columns in monkey striate cortex. *Philos. Trans. R. Soc. Lond. B: Biol. Sci.* 278, 377–409
- 39 Holtmaat, A. *et al.* (2009) Long-term, high-resolution imaging in the mouse neocortex through a chronic cranial window. *Nat. Protoc.* 4, 1128–1144
- 40 Chen, J.L. *et al.* (2011) Structural basis for the role of inhibition in facilitating adult brain plasticity. *Nat. Neurosci.* 14, 587–594
- 41 Vaughn, J.E. (1989) Fine structure of synaptogenesis in the vertebrate central nervous system. *Synapse* 3, 255–285
- 42 Lichtman, J.W. and Colman, H. (2000) Synapse elimination and indelible memory. *Neuron* 25, 269–278
- 43 Alsina, B. *et al.* (2001) Visualizing synapse formation in arborizing optic axons in vivo: dynamics and modulation by BDNF. *Nat. Neurosci.* 4, 1093–1101
- 44 Chen, A.C. *et al.* (2001) ECS-induced mossy fiber sprouting and BDNF expression are attenuated by ketamine pretreatment. *J. ECT* 17, 27–32
- 45 Bessa, J.M. *et al.* (2009) The mood-improving actions of antidepressants do not depend on neurogenesis but are associated with neuronal remodeling. *Mol. Psychiatry* 14, 764–773
- 46 Huttenlocher, P.R. and Dabholkar, A.S. (1997) Regional differences in synaptogenesis in human cerebral cortex. *J. Comp. Neurol.* 387, 167–178
- 47 Petanjek, Z. *et al.* (2011) Extraordinary neoteny of synaptic spines in the human prefrontal cortex. *Proc. Natl. Acad. Sci. U.S.A.* 108, 13281–13286
- 48 Maya Vetencourt, J.F. *et al.* (2008) The antidepressant fluoxetine restores plasticity in the adult visual cortex. *Science* 320, 385–388
- 49 Trachtenberg, J.T. *et al.* (2002) Long-term in vivo imaging of experience-dependent synaptic plasticity in adult cortex. *Nature* 420, 788–794
- 50 Chen, F. *et al.* (2009) Repeated electroconvulsive seizures increase the total number of synapses in adult male rat hippocampus. *Eur. Neuropsychopharmacol.* 19, 329–338
- 51 Chen, F. *et al.* (2008) Changes in rat hippocampal CA1 synapses following imipramine treatment. *Hippocampus* 18, 631–639
- 52 Hajszan, T. *et al.* (2009) Remodeling of hippocampal spine synapses in the rat learned helplessness model of depression. *Biol. Psychiatry* 65, 392–400
- 53 Hajszan, T. *et al.* (2005) Short-term treatment with the antidepressant fluoxetine triggers pyramidal dendritic spine formation in rat hippocampus. *Eur. J. Neurosci.* 21, 1299–1303
- 54 Cerqueira, J.J. *et al.* (2007) The prefrontal cortex as a key target of the maladaptive response to stress. *J. Neurosci.* 27, 2781–2787
- 55 Hajszan, T. *et al.* (2010) Effects of estradiol on learned helplessness and associated remodeling of hippocampal spine synapses in female rats. *Biol. Psychiatry* 67, 168–174
- 56 Radley, J.J. *et al.* (2006) Repeated stress induces dendritic spine loss in the rat medial prefrontal cortex. *Cereb. Cortex* 16, 313–320
- 57 Stewart, C.A. and Reid, I.C. (2000) Repeated ECS and fluoxetine administration have equivalent effects on hippocampal synaptic plasticity. *Psychopharmacology (Berl.)* 148, 217–223
- 58 Kobayashi, K. *et al.* (2010) Reversal of hippocampal neuronal maturation by serotonergic antidepressants. *Proc. Natl. Acad. Sci. U.S.A.* 107, 8434–8439
- 59 Sale, A. *et al.* (2007) Environmental enrichment in adulthood promotes amblyopia recovery through a reduction of intracortical inhibition. *Nat. Neurosci.* 10, 679–681
- 60 Karpova, N.N. *et al.* (2011) Fear erasure in mice requires synergy between antidepressant drugs and extinction training. *Science* 334, 1731–1734
- 61 Kirkwood, A. *et al.* (1995) Co-regulation of long-term potentiation and experience-dependent synaptic plasticity in visual cortex by age and experience. *Nature* 375, 328–331
- 62 Sale, A. *et al.* (2009) Enrich the environment to empower the brain. *Trends Neurosci.* 32, 233–239
- 63 Weikum, W.M. *et al.* (2012) Prenatal exposure to antidepressants and depressed maternal mood alter trajectory of infant speech perception. *Proc. Natl. Acad. Sci. U.S.A.* 109 (Suppl. 2), 17221–17227
- 64 Holderbach, R. *et al.* (2007) Enhanced long-term synaptic depression in an animal model of depression. *Biol. Psychiatry* 62, 92–100
- 65 West, A.E. and Greenberg, M.E. (2011) Neuronal activity-regulated gene transcription in synapse development and cognitive function. *Cold Spring Harb. Perspect. Biol.* 3, <http://dx.doi.org/10.1101/cshperspect.a005744>
- 66 Flavell, S.W. and Greenberg, M.E. (2008) Signaling mechanisms linking neuronal activity to gene expression and plasticity of the nervous system. *Annu. Rev. Neurosci.* 31, 563–590
- 67 Castrén, E. *et al.* (2012) Treatment of neurodevelopmental disorders in adulthood. *J. Neurosci.* 32, 14074–14079
- 68 Borrelli, E. *et al.* (2008) Decoding the epigenetic language of neuronal plasticity. *Neuron* 60, 961–974
- 69 Krishnan, V. and Nestler, E.J. (2008) The molecular neurobiology of depression. *Nature* 455, 894–902
- 70 Jiang, Y. *et al.* (2008) Epigenetics in the nervous system. *J. Neurosci.* 28, 11753–11759
- 71 Tsankova, N. *et al.* (2007) Epigenetic regulation in psychiatric disorders. *Nat. Rev. Neurosci.* 8, 355–367
- 72 Duman, R.S. and Newton, S.S. (2007) Epigenetic marking and neuronal plasticity. *Biol. Psychiatry* 62, 1–3
- 73 Guo, J.U. *et al.* (2011) Neuronal activity modifies the DNA methylation landscape in the adult brain. *Nat. Neurosci.* 14, 1345–1351
- 74 Poo, M.M. (2001) Neurotrophins as synaptic modulators. *Nat. Rev. Neurosci.* 2, 24–32
- 75 Bergami, M. *et al.* (2008) Deletion of TrkB in adult progenitors alters newborn neuron integration into hippocampal circuits and increases anxiety-like behavior. *Proc. Natl. Acad. Sci. U.S.A.* 105, 15570–15575
- 76 Li, Y. *et al.* (2008) TrkB regulates hippocampal neurogenesis and governs sensitivity to antidepressive treatment. *Neuron* 59, 399–412
- 77 McAllister, A.K. *et al.* (1999) Neurotrophins and synaptic plasticity. *Annu. Rev. Neurosci.* 22, 295–318



- 78 Kang, H. and Schuman, E.M. (1995) Long-lasting neurotrophin-induced enhancement of synaptic transmission in the adult hippocampus. *Science* 267, 1658–1662
- 79 Lu, B. (2003) BDNF and activity-dependent synaptic modulation. *Learn. Mem.* 10, 86–98
- 80 Saarelainen, T. *et al.* (2003) Activation of the TrkB neurotrophin receptor is induced by antidepressant drugs and is required for antidepressant-induced behavioral effects. *J. Neurosci.* 23, 349–357
- 81 Rossi, C. *et al.* (2006) Brain-derived neurotrophic factor (BDNF) is required for the enhancement of hippocampal neurogenesis following environmental enrichment. *Eur. J. Neurosci.* 24, 1850–1856
- 82 Berton, O. *et al.* (2006) Essential role of BDNF in the mesolimbic dopamine pathway in social defeat stress. *Science* 311, 864–868
- 83 Alter, M.D. *et al.* (2008) Variation in the large-scale organization of gene expression levels in the hippocampus relates to stable epigenetic variability in behavior. *PLoS ONE* 3, e3344
- 84 Alter, M.D. and Hen, R. (2009) Is there a genomic tone? Implications for understanding development, adaptation and treatment. *Dev. Neurosci.* 31, 351–357
- 85 Feinberg, A.P. and Irizarry, R.A. (2010) Stochastic epigenetic variation as a driving force of development, evolutionary adaptation, and disease. *Proc. Natl. Acad. Sci. U.S.A.* 107 (Suppl. 1), 1757–1764
- 86 Berardi, N. *et al.* (2003) Molecular basis of plasticity in the visual cortex. *Trends Neurosci.* 26, 369–378
- 87 Mountcastle, V.B. (1997) The columnar organization of the neocortex. *Brain* 120, 701–722
- 88 Maya Vetencourt, J.F. *et al.* (2011) Serotonin triggers a transient epigenetic mechanism that reinstates adult visual cortex plasticity in rats. *Eur. J. Neurosci.* 33, 49–57
- 89 Morishita, H. and Hensch, T.K. (2008) Critical period revisited: impact on vision. *Curr. Opin. Neurobiol.* 18, 101–107
- 90 Chollet, F. *et al.* (2011) Fluoxetine for motor recovery after acute ischaemic stroke (FLAME): a randomised placebo-controlled trial. *Lancet Neurol.* 10, 123–130
- 91 Aboukhatwa, M. *et al.* (2010) Antidepressants are a rational complementary therapy for the treatment of Alzheimer's disease. *Mol. Neurodegener.* 5, 10
- 92 Chadwick, W. *et al.* (2011) Amitriptyline-mediated cognitive enhancement in aged 3xTg Alzheimer's disease mice is associated with neurogenesis and neurotrophic activity. *PLoS ONE* 6, e21660
- 93 Han, X. *et al.* (2011) Imipramine treatment improves cognitive outcome associated with enhanced hippocampal neurogenesis after traumatic brain injury in mice. *J. Neurotrauma* 28, 995–1007